arXiv:quant-ph/0512225v1 24 Dec 2005

Husimi distribution function and one-dimensional Ising model

F. Kheirandish[1] *

[1] Department of Physics, University of Isfahan,
Hezar Jarib Ave., Isfahan, Iran.

Abstract

Husimi distribution function for the one-dimensional Ising model is obtained. One-point and joint distribution functions are calculated and their thermal behaviour are discussed.

PACS numbers: 02.70.-c, 03.65.Ca

1 Introduction

Husimi distribution function [1], is a positive definite distribution over the parameter space of a definite set of coherent states [2]. Positive definitness of the Husimi distribution function, makes it as a good candidate for investigating quantum features using a classical like distribution over the phase space. In this case, to each coherent state $|z\rangle$ of the system there corresponds a unique distribution function $\mu(z) = \frac{1}{Z}\langle z|e^{-\beta\hat{H}}|z\rangle$, where Z, is the partition function of the system. $\hat{H}$ and β are Hamiltonian and inverse temperature parameter respectively. In this paper, using the single-fermion or spin-$\frac{1}{2}$ coherent states, the coherent states of a one-dimensional spin chain are obtained. The Husimi distribution function is then obtained for the one-dimensional Ising model as the prototype of some important models in statistical mechanics. One-point and joint distribution functions or correlation functions are calculated and their thermal behaviour is discussed.

*fardin_kh@phys.ui.ac.ir

2 One-dimensional Ising model

Hamiltonian of the one dimensional Ising model in a homogeneous magnetic field B along the z-axis is

$$\hat{H} = -J \sum_{k=1}^{N} \hat{S}_{k,z} \hat{S}_{k+1,z} + B \sum_{k=1}^{N} \hat{S}_{k,z}, \tag{1}$$

where periodic boundary condition is assumed, i.e, $N+1 \equiv 1$. The Hilbert space $\aleph$ of this model is a 2^N-dimensional vector space spaned by the following tensor products as basis vectors

$$|\vec{i}\rangle = |i_1\rangle \otimes |i_2\rangle \otimes ... \otimes |i_N\rangle = \prod_{k=1}^{N} \otimes |i_k\rangle, \tag{2}$$

where $|i_k\rangle \in \{|+\rangle, |-\rangle\}$, for $k = 1...N$. The basis vectors $|\vec{i}\rangle$, are Hamiltonian eigenvectors with respective eigenvalues $E_{\vec{i}}$,

$$\begin{aligned} \hat{H}|\vec{i}\rangle &= (-J \sum_{k=1}^{N} i_k i_{k+1} + B \sum_{k=1}^{N} i_k)|\vec{i}\rangle, \\ E_{\vec{i}} &= -J \sum_{k=1}^{N} i_k i_{k+1} + B \sum_{k=1}^{N} i_k, \end{aligned} \tag{3}$$

3 Ising model coherent states

Coherent states of the one dimensional Ising model can be defined from the tensor product of the single-fermion coherent states [2]. A single-fermion coherent state for the spin half ($s = \frac{1}{2}$) representation can be written like this

$$|z\rangle = \sin(\frac{\theta}{2}) \exp(-i\varphi)|+\rangle + \cos(\frac{\theta}{2})|-\rangle, \tag{4}$$

where $z = (\frac{\theta}{2}) exp(-i\varphi)$, $0 \leq \theta \leq \pi$, $0 \leq \varphi \leq 2\pi$ and $|+\rangle$, $|-\rangle$ are $\hat{S}_z$ eigenvectors. The completeness of the fermion coherent states is

$$\frac{1}{2\pi} \int d\Omega |z\rangle\langle z| = 1, \tag{5}$$

where $d\Omega = \sin\theta d\theta d\varphi$. Tensor product of these fermion coherent states makes Ising model coherent states

$$|\vec{z}\rangle = \prod_{k=1}^{N} \otimes (\sin(\frac{\theta_k}{2}) \exp(-i\varphi_k)|+\rangle + \cos(\frac{\theta_k}{2})|-\rangle), \tag{6}$$

the scalar product between a coherent state $|\vec{z}>$ and an eigenvector $|\vec{i}>$, of the Hamiltonian $\hat{H}$, is

$$\langle\vec{z}|\vec{i}\rangle = \prod_{k=1}^{N}(\sin(\frac{\theta_k}{2})\exp(-i\varphi_k)\delta_{i_k,1} + \cos(\frac{\theta_k}{2})\delta_{i_k,-1}), \tag{7}$$

so

$$|\langle\vec{z}|\vec{i}\rangle|^2 = \prod_{k=1}^{N}(\sin^2(\frac{\theta_k}{2})\delta_{i_k,1} + \cos^2(\frac{\theta_k}{2})\delta_{i_k,-1}). \tag{8}$$

Partition function of the Hamiltonian (1) is defined as

$$Z = \sum_{\vec{i}}\langle\vec{i} \mid \exp(-\beta\hat{H}) \mid \vec{i}\rangle = tr(\hat{T}^N), \tag{9}$$

where

$$\hat{T} = \begin{pmatrix} \exp(\beta(J-B)) & \exp(-\beta J) \\ \exp(-\beta J) & \exp(\beta(J+B)) \end{pmatrix}, \tag{10}$$

is the transition matrix.

4 Husimi distribution function

Husimi distribution function is a normalized positive definite distribution in the parameter space of coherent states and is defined as follows

$$\begin{aligned} \mu(\vec{z}) &= \frac{1}{Z}\langle\vec{z}|\exp(-\beta\hat{H})|\vec{z}>, \\ &= \frac{1}{Z}\sum_{\{\vec{i}\}}\langle\vec{z}|\exp(-\beta\hat{H})|\vec{i}\rangle\langle\vec{i}|\vec{z}>, \\ &= \frac{1}{Z}\sum_{\{\vec{i}\}} e^{\beta J\sum_{k=1}^{N} i_k i_{k+1} - \beta B\sum_{k=1}^{N} i_k}|\langle\vec{z}|\vec{i}\rangle|^2, \end{aligned} \tag{11}$$

substituting (8) in (11) and doing some routine calculations, we get the following relation for Husimi distribution

$$\begin{aligned} \mu(\vec{z}) &= \frac{1}{2^N}\{1 - \sum_{m=1}^{N}\langle\hat{S}_{m,z}\rangle u_m + \sum_{m<n}\langle\hat{S}_{m,z}\hat{S}_{n,z}\rangle u_m u_n + \cdots \\ &+ (-1)^N\langle\hat{S}_{1,z}\hat{S}_{2,z}\cdots\hat{S}_{N,z}\rangle u_1 u_2 \cdots u_n\}, \end{aligned} \tag{12}$$

where for simplicity $u_k = cos\theta_k$ is assumed and a general n-point corelation function $\langle \hat{S}_{m,z}\hat{S}_{n,z}\cdots\hat{S}_{r,z}\rangle$ is defined as

$$\langle \hat{S}_{m,z}\hat{S}_{n,z}\cdots\hat{S}_{r,z}\rangle = \frac{1}{Z}\sum_{\vec{i}} i_m i_n \cdots i_r e^{\beta J\sum_{k=1}^{N} i_k i_{k+1} - \beta B \sum_{k=1}^{N} i_k}, \qquad (13)$$

where $\hat{S}_{k,z}|i_k\rangle = i_k|i_k\rangle$, from (12) it is clear that the Husimi distribution function is independent of parameters φ_k and contains the information about all the n-point correlation functions, i.e., it is the generator of the n-point functions of the Ising model

$$\begin{aligned}\langle \hat{S}_{r_1,z}\hat{S}_{r_2,z}\cdots\hat{S}_{r_m,z}\rangle &= (-3)^m \int\cdots\int \mu(\vec{z}) u_{r_1}u_{r_2}\cdots u_{r_m}\prod_{k=1}^{N} du_k, \\ &= (-3)^m \langle u_{r_1}u_{r_2}\cdots u_{r_m}\rangle. \qquad (14)\end{aligned}$$

Now the Husimi distribution function related to the cite k, can be calculated by integrating over parameters of the phase spase except those belong to the kth cite, so

$$\mu(u_k) = \frac{1}{2}(1 - u_k \frac{tr[\hat{T}^N\sigma_z]}{tr[\hat{T}^N]}), \qquad (15)$$

where as usual σ_z is

$$\sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \qquad (16)$$

The joint Husimi distribution function related to cites m and n, is equal to

$$\mu(u_m, u_n) = \frac{1}{4}(1 - (u_m + u_n)\frac{tr[\hat{T}^N\sigma_z]}{tr[\hat{T}^N]} + u_m u_n \frac{tr[\hat{T}^{N-n+m}\sigma_z\hat{T}^{n-m}\sigma_z]}{tr[\hat{T}^N]}), \quad (17)$$

for calculating $tr[\hat{T}\sigma_z]$ and $tr[\hat{T}^{N-n+m}\sigma_z\hat{T}^{n-m}\sigma_z]$ we can diagonalize the matrix $\hat{T}$, and write the matrix σ_z in the basis of eigenvectors of $\hat{T}$, the eigenvalues of $\hat{T}$ are

$$\lambda_\pm = \exp(\beta J)\cosh(\beta B) \pm \sqrt{\exp(2\beta J)\cosh^2(\beta B) - 2\sinh(\beta J)}, \qquad (18)$$

and the transformation matrix between the standard basis $\{|+\rangle, |-\rangle\}$ and the eigenvectors of $\hat{T}$ is

$$U = \begin{pmatrix} \sin(\omega) & -\cos(\omega) \\ \cos(\omega) & \sin(\omega) \end{pmatrix}, \qquad (19)$$

where $|\lambda_+\rangle = \begin{pmatrix} sin(\omega) \\ cos(\omega) \end{pmatrix}$ and $|\lambda_-\rangle = \begin{pmatrix} -cos(\omega) \\ sin(\omega) \end{pmatrix}$ are normalized eigenvectors of $\hat{T}$ belonging to eigenvalues λ_+ and λ_- respectively and

$$\tan(\omega) = \frac{\exp(-\beta J)}{\exp(\beta J)\sinh(\beta B) + \sqrt{\exp(2\beta J)\sinh^2(\beta B) + \exp(-2\beta J)}}. \quad (20)$$

Now in the basis $|\lambda_+\rangle$, $|\lambda_-\rangle$, we have

$$\begin{aligned} tr[\hat{T}\sigma_z] &= tr(\begin{pmatrix} \lambda_+^N & 0 \\ 0 & \lambda_-^N \end{pmatrix} \begin{pmatrix} -\cos(2\omega) & -\sin(2\omega) \\ -\sin(2\omega) & \cos(2\omega) \end{pmatrix}), \\ &= tr[\begin{pmatrix} -\lambda_+^N \cos(2\omega) & -\lambda_+^N \sin(2\omega) \\ -\lambda_-^N \sin(2\omega) & \lambda_-^N \cos(2\omega) \end{pmatrix}], \\ &= (\lambda_-^N - \lambda_+^N)\cos(2\omega), \end{aligned} \quad (21)$$

and similarly

$$\begin{aligned} tr[\hat{T}^{N-j+i}\sigma_z\hat{T}^{j-i}\sigma_z] &= \lambda_+^N \cos^2(2\omega) + \lambda_+^{N-j+i}\lambda_-^{j-i}\sin^2(2\omega) \\ &+ \lambda_-^{N-j+i}\lambda_+^{j-i}\sin^2(2\omega) + \lambda_-^N\cos^2(2\omega), \end{aligned} \quad (22)$$

substituting (21) and (22) in (15) and (17) respectively, we obtain

$$\mu(u_k) = \frac{1}{2}(1 - \cos(2\omega)[\frac{\lambda_-^N - \lambda_+^N}{\lambda_-^N + \lambda_+^N}]u_k), \quad (23)$$

$$\begin{aligned} \mu(u_i, u_j) &= \frac{1}{4}(1 - (u_i + u_j)\cos(2\omega)[\frac{\lambda_-^N - \lambda_+^N}{\lambda_-^N + \lambda_+^N}]u_k + u_i u_j \cos^2(2\omega) \\ &+ u_i u_j \sin^2(2\omega)\frac{\lambda_+^{N-j+i}\lambda_-^{j-i} + \lambda_-^{N-j+i}\lambda_+^{j-i}}{\lambda_-^N + \lambda_+^N}), \end{aligned} \quad (24)$$

in thermodynamic limit, i.e., $N \to \infty$, we have

$$\begin{aligned} \mu(u_k) &= \frac{1}{2}(1 + \cos(2\omega)u_k), \\ \mu(u_i, u_j) &= \frac{1}{4}(1 + (u_i + u_j)\cos(2\omega) + u_i u_j \cos^2(2\omega) \\ &+ u_i u_j (\frac{\lambda_-}{\lambda_+})^{j-i}\sin^2(2\omega)), \end{aligned} \quad (25)$$

such that

$$\begin{aligned}
\cos(2\omega) &= \frac{2\exp(\beta J)\sinh(\beta B)}{\sqrt{4\exp(2\beta J)\sinh^2(\beta B)+4\exp(-2\beta J)}}, \\
&= \frac{Sgn(B)}{\sqrt{1+\frac{\exp(-4\beta J)}{\sinh^2(\beta B)}}}, \\
\tan(2\omega) &= \frac{exp(-2\beta J)}{\sinh(\beta B)},
\end{aligned} \tag{26}$$

where $Sgn(B)$, is the sign function. The behaviour of one-point distribution function $\mu(u_k)$, in high and low temperatures, is

$$\mu(u_k) = \begin{cases} \frac{1}{2} & \beta \to 0, \\ \frac{1}{2}(1+u_k) & \beta \to \infty, \end{cases} \tag{27}$$

in zero temperature, $\mu(u_k)$ attains it's maximum for $\theta_k = 0$, which from (4) corresponds to the spin down state, $|-\rangle$, as expected.
Similarly we can find the following behaviour for joint distribution $\mu(u_i, u_j)$,

$$\mu(u_i, u_j) = \begin{cases} \frac{1}{4} & \beta \to 0, \\ \frac{1}{4}(1+u_i+u_j+u_iu_j) & \beta \to \infty, \end{cases} \tag{28}$$

in this case the disjoint distribution attains it's maximum value for $\theta_i = \theta_j = 0$, which means that most probably the two spins are in the spin down state, i.e., $|-\rangle$, in derivation of these behaviours it is assumed that $Sign(B) = 1$, by reversing the magnetic field ($Sign(B) = -1$), the spins flip to the spin up state.

References

[1] K. Husimi, Proc. Phys. Math. Soc. Jpn. 22, 264 (1940)

[2] W. M. Zhang, D. H. Feng and R. Gilmore, Rev. Mod. Phys. Vol.62, No.4, 867 (1990)